Cronartium flaccidum references:

Alexopoulos CJ, Mims CW, Blackwell M, 1996. Introductory Mycology. Fourth edition. New York, New York, USA: John Wiley and Sons, Inc, 868 pp.

Azbukina, ZM, 2008. Taxonomical notes on species of *Cronartium* (Uredinales) occurring in Russia. Mikologiya i Fitopatologiya, 42:3-12.

BPI (US National Fungus Collections), 2009. Fungal Databases – Specimens. Systematic Mycology and Microbiology Laboratory/Agricultural Research Service/USDA. Beltsville, Maryland, USA. www.nt.ars-grin.gov/fungaldatabases/specimens/specimens.cfm. Accessed June 18, 2009.

Butin H, 1995. Tree Diseases and Disorders. Oxford, United Kingdom: Oxford University Press, 252 pp.

CABI, 1989. *Cronartium flaccidum* Commonwealth Mycological Institute Distribution Maps of Plant Diseases, 616:1-2.

Canadian Food Inspection Agency, 2008. Wood packaging. Ottawa, Canada: Government of Canada. http://www.inspection.gc.ca/english/plaveg/for/cwpc/wdpkge.shtml Accessed July 2, 2009.

Cao ZM, Li ZQ, Zhuang JY, 2000. Uredinales from the Qinling Mountains. Mycosystema, 19:13-23.

Chen MM, 2002. Forest Fungi Phytogeography: Forest Fungi Phytogeography of China, North America, and Siberia and International Quarantine of Tree Pathogens. Sacramento, California: Pacific Mushroom Research and Education Center, 495 pp.

Cho WD, Shin HD, eds., 2004. List of plant diseases in Korea. Fourth edition. Seoul, Republic of Korea: Korean Society of Plant Pathology, 779 pp.

Denchev CM, 1995. Bulgarian Uredinales. Mycotaxon, 55:405-465.

Dudka IO, Heluta VP, Tykhonenko YY, Andrianova TV, Hayova VP, Prydiuk MP, Dzhagan VV, Isikov VP, 2004. Fungi of the Crimean Peninsula. Kiev, Ukraine: M.G. Kholodny Institute of Botany, National Academy of Sciences of Ukraine, 452 pp.

Gibbs J, 1997. Resin top disease. In: Compendium of conifer diseases. Hansen EM, Lewis KJ, eds. St. Paul, Minnesota, USA: American Phytopathological Society, 30-31.

Gibbs JN, England N, Wolstenholme R, 1988. Variation in the pine stem rust fungus *Peridermium pini* in the United Kingdom. Plant Pathology, 37:45–53.

Gibbs JN, Grieg BJW, Hickman T, 1987. An analysis of *Peridermium* stem rust of Scots pine in Thetland Forest in 1984-1985. Forestry, 60:203-218.

Gjaerum, H. 1974. Nordens

Rustsopper. Fungiflora, Oslo,

321 pp.

Goncalves da Cunha A, 1936. Uredineas de Portugal. Boletin da Sociedade Broteriana, 11:169-265.

Greig BJW, 1987. History of *Peridermium* stem rust of Scots pine (*Pinus sylvestris* L.) in Thetford Forest, East Anglia. Forestry, 60:193-202.

Hantula J, Kasanen R, Kaitera J, Moricca S, 2002. Analyses of genetic variation suggest that pine rusts *Cronartium flaccidum* and *Peridermium pini* belong to the same species. Mycological Research, 106:203-209.

Hantula J, Maarit Niemi E, Kaitera J, Jalkanen R, Kurkela T, 1998. Genetic variation of the resin top fungus in Finland as determined by random amplified microsatellites (RAMS). European Journal of Forest Pathology, 28:361-372.

Hiratsuka Y, 1968. Morphology and cytology of aeciospores and aeciospore germ tubes of host-alternating and pine-to-pine races of *Cronartium flaccidum* in northern Europe. Canadian Journal of Botany, 46:1119-1122.

Hiratsuka Y, 1969. *Endocronartium*, a new genus for autoecious pine stem rusts. Canadian Journal of Botany, 47:1493-1495.

Hiratsuka N, Chen ZC, 1991. A list of Uredinales collected from Taiwan. Transactions of the Mycological Society of Japan, 32:3-22.

Hunt R, 1997. Stem rusts. In: Compendium of conifer diseases. Hansen EM, Lewis KJ, eds. St. Paul, Minnesota, USA: American Phytopathological Society, 26.

Kaitera J, 1999. *Cronartium flaccidum* fruitbody production on *Melampyrum* spp. and some important alternate hosts to pine. European Journal of Forest Pathology, 29:391–398.

Kaitera J, 2000. Analysis of *Cronartium flaccidum* lesion development on pole-stage Scots pines. Silva Fennica, 34:21–27.

Kaitera J, Aalto T, Jalkanen R, 1994. Effect of resin-top disease caused by *Peridermium pini* on the volume and value of *Pinus sylvestris* saw timber and pulpwood. Scandinavian Journal of Forest Research, 9:376–381.

Kaitera J, Hantula J, 1998. *Melampyrum sylvaticum*, a new alternate host for pine stem rust *Cronartium flaccidum*. Mycologia, 90:1028-1030.

Kaitera J, Seitamake L, Hantula J, Jalkanen R, Kurkela T, 1999a. Inoculation of known and potential alternate hosts with *Peridermium pini* and *Cronartium flaccidum* aeciospores. Mycological Research, 103:235-241.

Kaitera J, Seitamake L, Hantula J, Jalkanen R, Kurkela T, 1999b. Morphological variation of *Peridermium pini* and *Cronartium flaccidum* aeciospores. Mycological Research, 103:677–683.

Kaitera J, Nuorteva H, 2006. Finnish *Cronartium ribicola* does not infect alternate hosts of *Cronartium flaccidum.* Forest Pathology, 36:225-246.

Kaitera J, Nuorteva H, 2008. Inoculations of eight *Pinus* species with *Cronartium* and *Peridermium* stem rusts. Forest Ecology and Management, 25:973-981.

Kaitera J, Nuorteva H, Hantula J, 2005. Distribution and frequency of *Cronartium flaccidum* on *Melampyrum* spp. in Finland. Canadian Journal of Forest Research, 35:229-234.

Kasanen R, 1997. Aeciospores of *Cronartium flaccidum, C. ribicola* and *Endocronartium pini* show no differences in morphology. European Journal of Forest Pathology, 27:251-260.

Kasanen R, Kaitera J, Hantula J, 2000. The genetic structure of *Peridermium pini and Cronartium flaccidum* cankers on Scots pine as revealed by two multi-allelic loci. Forest Pathology, 30:221–230.

Kobayashi T, 2007. Index of Fungi Inhabiting Woody Plants in Japan. Host, Distribution and Literature. Tokyo, Japan: Zenkoku-Noson-Kyiku Kyokai Publishing Co., Ltd., 1227 pp.

Kuprevich VF, Tranzschel VG, 1957. Rust fungi. I. Family Melampsoraceae. In: Cryptogamic Plants of the USSR. Vol. 4. Savich VP, ed. Moscow – Leningrad: Academy of Sciences of the USSR, Komarov Institute of Botany, 518 pp. (translated by the Israel Program for Scientific Translations, Jerusalem, 1970).

Kuzmina NA, Kuz'min SR, 2008. Intraspecific response of Scots pine (*Pinus sylvestris* L.) to pathogens in a provenance trial in middle Siberia. Eurasian Journal of Forest Research, 11:51-59.

MAF, 2009. Search results – *Endocronartium pini*. Biosecurity New Zealand, Ministry of Agriculture and Forestry. http://www.biosecurity.govt.nz/search/node/Endocronartium+pini. Accessed July 2, 2009.

Martinsson O, Nilsson B, 1987. The impact of *Cronartium flaccidum* on the growth of *Pinus sylvestris*, Scandinavian Journal of Forest Research, 2:349–357.

Mittempergher L, Raddi P, 1977. Variation of diverse sources of *Cronartium flaccidum*. European Journal of Forest Pathology, 7:93-98.

Mordue JEM, Gibson IAS, 1978. *Cronartium flaccidum*. CMI Descriptions of Phytopathogenic Fungi and Bacteria, 580:1-2.

Moricca S, Kasuga T, Mitchelson K, Ragazzi A, Diamandis S, 1996. Heterogeneity in intergenic regions of the ribosomal repeat of the pine-blister rusts *Cronartium flaccidum* and *Peridermium pini*. Current Genetics, 29:388-394.

Moricca S, Ragazzi A, 1996. Culture characteristics and variation of *Cronartium flaccidum* isolates. Canadian Journal of Botany, 74:924-933.

Moricca S, Ragazzi A, 1998. Use of RFLP and SSCP analysis to differentiate the pine rusts *Cronartium flaccidum and Peridermium pini*. Mycological Research, 102:666–670.

Moricca S, Ragazzi A, Mitchelson KR, Assante G, 2001. Antagonism of the two-needle pine stem rust fungi *Cronartium flaccidum* and *Peridermium pini* by *Cladosporium tenuissimum* in vitro and in planta. Phytopathology, 91:457-68.

Moricca S, Ragazzi A, 2008. Biological and integrated means to control rust diseases. In: Integrated Management of Plant Pests and Diseases. Ciancio A, Mukerji KG, eds. Vol. 3. Dordrecht, The Netherlands: Springer, 303-329.

Mulenko W, Kozlowska M, Salata B, 2004. Microfungi of the Tatra National Park. A checklist. W. Szafer Institute of Botany, Polish Academy of Sciences, 72 pp.

NCBI, 2009. Entrez cross-database search engine. National Center for Biotechnology Information, U.S. National Library of Medicine, Bethesda, Maryland, USA. http://www.ncbi.nlm.nih.gov/sites/gquery. Accessed June 29, 2009.

Neumann FG, Marks GC, 1990. Status and management of insect pests and diseases in Victorian softwood plantations. Australian Forestry, 53:131- 144.

Pappinen A, von Weissenberg K, 1994. The ability of the pine-top weevil to carry spores and infect Scots pine with *Endocronartium pini*. European Journal of Forest Pathology, 24:258-263.

Pei MH, Gibbs JN, 1991. Cultural variation in *Peridermium pini* from two regions of the United Kingdom: Scotland and East Anglia. European Journal of Forest Pathology, 21:164-167.

Raddi P, Fagnini A, 1978. Relative susceptibility to blister rust caused by *Cronartium flaccidum* of several species of pine. European Journal of Forest Pathology, 8:58-61.

Raddi P, Mittempergher L, Moriondo F, 1979. Testing of *Pinus pinea* and *P. pinaster* for resistance to *Cronartium flaccidum*. Phytopathology, 69:679- 681.

Ragazzi A, 1983. Development of *Cronartium flaccidum* (Alb. et Schw.) Wint. on *Vincetoxicum officinale* Moench in connection with some environmental factors. Journal of Phytopathology, 108:160-171.

Ragazzi A, 1989. Past, present and future of *Cronartium flaccidum* in Italy. Phytopathologia Mediterranea, 28:5–9.

Ragazzi A, Dellavalle Fedi I, Mesturino L, 1986. *Cronartium flaccidum* on *Pinus* spp.: relation of inoculum concentration to symptom development. European Journal of Forest Pathology, 16:16-21.

Ragazzi A, Fagnani A, Dellavalle Fedi I, 1987. Telial and basidiospore stages of *Cronartium flaccidum*: light and scanning electron microscope observations. Phytopathologia Mediterranea, 26:113-116.

Roll-Hansen F, 1973. Resistance of *Paeonia* cultivars to *Cronartium flaccidum* in Norway. European Journal of Forest Pathology, 3:142-145

Rossman AY, Britton K, Luster D, Palm M, Royer MH, Sherald J, 2006. Evaluating the threat posed by fungi on the APHIS list of regulated plant pests. Plant Health Progress. (Online.) http://www.plantmanagementnetwork.org/php/

Sinclair WA, Lyon H, 2005. Diseases of Trees and Shrubs. Second edition. Ithaca, New York: Cornell University Press. 660 pp.

Smith IM, Dunez J, Lelliott RA, Phillips DH, Archer SA, 1988. European Handbook of Plant Diseases. Oxford, UK: Blackwell Scientific Publications, 583 pp.

Tai FL, 1979. Sylloge Fungorum Sinicorum. Beijing, China: Science Press Academica Sininica, 1527 pp.

Teng SC, 1996. Fungi of China. Ithaca, New York, USA: Mycotaxon Ltd, 663 pp.

USDA/APHIS, 2008. Circular 34: Diseases of *Picea* spp. Post-entry Quarantine Manual for State Inspectors. Washington, D.C., USA, pp. 10-151-153. http://www.aphis.usda.gov/import_export/plants/manuals/domestic/downloads/postentry.pdf

USDA-ARS, 2009. Germplasm Resources Information Network (GRIN). Online Database. National Germplasm Resources Laboratory, Beltsville, Maryland, USA. http://www.ars-grin.gov/cgi-bin/npgs/html/tax_search.pl. Accessed June 21, 2009.

USDA-NRCS, 2008. The PLANTS Database. National Plant Data Center, Baton Rouge, Louisiana, USA. http://plants.usda.gov/ Accessed June 29, 2009.

Van der Kamp BJ, 1970. *Peridermium pini* (Pers.) Lév. and the resin-top disease of Scots pine. III. Infection and lesion development. Forestry, 43:73–88.

Vogler DR, Bruns TD, 1998. Phylogenetic relationships among the pine stem rust fungi (*Cronartium* and *Peridermium* spp.). Mycologia, 90:244-257.

Wilson M, Henderson DM, 1966. British Rust Fungi. Cambridge, United Kingdom: The Cambridge University Press, 384 +xvii pp.

Zhuang JY, Wei SX, 2005. Urediniomycetes, Uredinales. In: Fungi of Northwestern China. Zhuang WY, ed. Ithaca, New York, USA; Mycotaxon, Ltd., 430 pp.